


Sea current circulation around the Portofino Promontory










Now it is different: a great part of crew is made up of Romanian guys


First Regional Symposium on Sustainable small-scale [isheries in the Mediteranean and Black sea, 27-30 November 2013, St. Jualian's, Matra

GIORNO UE ACQUISTI DA ALBEISA	QUALITA	N. C3800	Kg. came	P1880	PREZZO	IMPORTO	Log-book of catches THE COOPERATIVE OF CAMOGLI
one li	BONITTO. LEPCLOSTE			3. 116	2.00 16,00	\mathcal{S}	FISHERMEN
<u>Sina</u>		 				<u> </u>	
DOMENIE A	19'L UG.	N (3912		PROV. PESO	PREZZO	IMPORTO	
ALBEITA One fi	BOULISTI PAIANITA SENNE TONNELLE AGONETH	3		6 18 1 1 2 2 10	£,00 6,00 4,00 4,00 4,00	X	
Seha	TONNEWA		Į	<u>{-145</u>	(, p.o.		
RIEPILOGO	Ven	uisti k dite k VI k			IOTALE E		









index was used

5

		1950-54 Balestra	1955-59	1960-64	1965-69	1970-74	1996-00	2000-05	2006-12
		et al., 1976	Balestra et al., 1976	Balestra et al., 1976	Balestra et al., 1976	Balestra et al., 1976	Relini 2001	this study	this study
Auxis rochei	avg (t)	21.39	11.15	7.03	4.69	4.12	9.50	8.13	5.78
Belone belone	avg (t)							0.47	0.87
Boops boops	avg (t)	4.02	2.96	1.08	1.40	2.92	4.00	3.42	1.54
Coryphaena hippurus	avg (t)							0.12	0.10
Euthynnus alletteratus	avg (t)							0.72	1.12
Oblada melanura	avg (t)	0.02	0.35	0.46	1.41	1.40		0.59	0.41
Sarda sarda	avg (t)	2.56	9.14	2.59	3.93	2.91	2.90	4.37	5.49
Sardinella sp.	avg (t)							0.23	0.40
Sarpa salpa	avg (t)	0.71	1.37	3.06	8.66	14.16	1.50	0.99	0.29
Scomber colias	avg (t)	2.49	2.21	3.22	0.41	2.56	2.40	4.50	7.11
Scomber scombrus	avg (t)	0.38	0.81	1.44	0.16	0.47	0.20	0.04	0.00
Scomberesox saurus	avg (t)	1.33	3.27	2.55	3.35	1.51	6.00	0.13	0.04
Seriola dumerili	avg (t)						9.00	9.72	4.84
Sphyraena viridensis	avg (t)							0.13	0.31
Trachurus spp.	avg (t)	3.47	2.12	5.53	1.66	7.05	12.00	17.24	6.88
Others	avg (t)	3.92	1.89	2.19	2.81	3.38	4.50	2.70	2.00
Total catches	avg (t)	40.43	34.49	29.71	32.02	41.46	52.00	53.28	37.18
	std	6.92	6.09	8.04	5.67	11.85	13.04	11.79	14.78

Average catches (t) from 1950 to 2012

First Regional Symposium on Sustainable small-scale fisheries in the Mediteranean and Black sea, 27-30 November 2013, St. Jualian's, Malta



Their catches in the season are related to warm seawater temperature (such as also Atlantic bonito) First Regional Symposium on Sustainable small-scale [isheries in the Mediteranean and Black Sea, 27-30 November 2013, St. Jualian's, J







29th May 2010: Associazione Culturale Ziguele

Aim of the cultural association:

the promotion and valorisation of the maritime culture and small scale fisheries tradition in the Mediterranean Sea, with particular regards to the sustainable fisheries





Since 2011

Slow Food Presidium

from the website www.presidislowfood.it:

The Camogli tonnarella is the only one remaining in Liguria and one of the few still active in all of Italy. The fishing system is highly sustainable because only medium and large fish remained trapped in the chambers. The mechanism does not immediately kill the fish, just concentrates them, so that the fishermen can check the catch

and release any fish that cannot be sold while they are still alive. The Presidio originates to promote and support fishermen activity because of the lack of interested people in this

hard activity



The Cooperative promotes guided tours to the Tonnarella in collaboration with a local no profit association







www.ziguele.it phone 3772900068 ziguele@gmail.com

