



Stock Assessment Form

Demersal species

Reference Year: 2016

Reporting Year: 2017

Trawl fishery data for the period 2004-2016 have been used to assess the *Mullus barbatus* stock in the GSA 06. The assessment has been carried out applying tuned VPA (Extended Survivor Analysis, XSA). FLR libraries under R language and Y/R analysis. Results from VPA indicate the average fishing mortality for ages 1-2 shows a general decreasing trend over the studied period reflecting the steady reduction observed in fishing effort. The fishery is in high overfishing with relative high biomass and spawning stock biomass.

Stock Assessment Form version 1.0 (January 2014)

Uploader: *Encarnacion Garcia Rodriguez*

Stock assessment form

Contents

1	Basic Identification Data.....	3
2	Stock identification and biological information	5
2.1	Stock unit.....	5
2.2	Growth and maturity.....	5
3	Fisheries information.....	7
3.1	Description of the fleet.....	7
3.2	Historical trends	8
3.3	Management regulations	8
3.4	Reference points.....	9
4	Fisheries independent information	10
4.1	{TYPE OF SURVEY}.....	10
4.1.1	Brief description of the direct method used	10
4.1.2	Spatial distribution of the resources	16
4.1.3	Historical trends	17
5	Ecological information.....	18
5.1	Protected species potentially affected by the fisheries	18
5.2	Environmental indexes	18
6	Stock Assessment	19
6.1	Tunnet VPA (XSA) (Darby & Flatman, 1994) and FLR libraries.	19
6.1.1	Model assumptions	19
6.1.2	Scripts.....	19
6.1.3	Input data and Parameters	19
6.1.4	Results	20
6.1.5	Robustness analysis.....	21
6.1.6	Retrospective analysis, comparison between model runs, sensitivity analysis, etc.	22
6.1.7	Assessment quality.....	23
6.2	Y/R (version 3.1.1; NOAA Fisheries Tools).....	23
6.3	STOCK / RECRUITMENT RELATIONSHIP	23
7	Stock predictions	24

7.1	Short term predictions	24
7.2	Medium term predictions.....	24
7.3	Long term predictions	24
8	Draft scientific advice	25
8.1	Explanation of codes	26

1 Basic Identification Data

Scientific name:	Common name:	ISCAAP Group:
Mullus barbatus	Red mullet	33 MUT[
1st Geographical sub-area:	2nd Geographical sub-area:	3rd Geographical sub-area:
[GSA_6]		
4th Geographical sub-area:	5th Geographical sub-area:	6th Geographical sub-area:
1st Country	2nd Country	3rd Country
Spain		
4th Country	5th Country	6th Country
Stock assessment method: (direct, indirect, combined, none)		
Indirect: XSA, FLR, YIELD PER RECRUIT		
Authors:		
García-Rodríguez, E. (1); Vivas, M. (1); Herrera, E. (1); Esteban, A. (1); Pérez-Gil, J. L. (2)		
Affiliation:		
(1) Instituto Español de Oceanografía (IEO). Murcia. Spain.		
(2) Instituto Español de Oceanografía (IEO). Málaga. Spain.		

The ISSCAAP code is assigned according to the FAO 'International Standard Statistical Classification for Aquatic Animals and Plants' (ISSCAAP) which divides commercial species into 50 groups on the basis of their taxonomic, ecological and economic characteristics. This can be provided by the GFCM secretariat if needed. A list of groups can be found here:

<http://www.fao.org/fishery/collection/asfis/en>

Direct methods (you can choose more than one):

- Acoustics survey
- Egg production survey
- Trawl survey
- SURBA

- Other (please specify)

Indirect method (you can choose more than one):

- ICA
- VPA
- LCA
- AMCI
- XSA
- Biomass models
- Length based models
- Other (please specify)

Combined method: you can choose both a direct and an indirect method and the name of the combined method (please specify)

2 Stock identification and biological information

Specify whether the assessment is considered to cover a complete stock unit. If the stock unit limits are more or less known, but for technical reasons the assessment only covers part of the stock (e.g. a GSA area but stock spreads to other GSAs), explain the state of the art of the stock unit knowledge. If there are doubts about the stock unit, state them here. If there is knowledge on migration rates between different stock units that affect the stock state them here.

2.1 Stock unit

Due to the lack of information about the structure of the population in the Western Mediterranean, it is considered that the stock limits of the assessed *Mullus barbatus* are in agreement with the limits of GSA 06.

2.2 Growth and maturity

Incorporate different tables if there are different maturity ogives (e.g. catch and survey). Also incorporate figures with the ogives if appropriate. Modify the table caption to identify the origin of the data (catches, survey). Incorporate names of spawning and nursery areas and maps if available.

Table 2.2-1: Maximum size, size at first maturity and size at recruitment.

Somatic magnitude measured (LT, LC, etc)				Units	
Sex	Fem	Mal	Combined	Reproduction season	May-July
Maximum size observed			30 (1)	Recruitment season	October-December
Size at first maturity			13.7 (2)	Spawning area	Continental shelf (4)
Recruitment size to the fishery			7.8 for the period 2004-2010 13.7 for the period 2011-2016 (3)	Nursery area	Coastal areas

(1) Size composition of trawl catches in GSA06.

(2) From the Spanish DCF National Programme (2016)

(3) García-Rodríguez, M. and Fernández, A. M. 2005.

(4) Lombarte, A.; L. Recasens; M. González and L. Gil de Sola (2000)

Table 2-2.4: M vector and proportion of matures by size or age (both sex)

Size/Age	Natural mortality	Proportion of matures
Age 0	1.23	0.127
Age 1	0.41	0.929
Age 2	0.28	0.999
Age 3+	0.22	1.000

Table 2-3: Growth and length weight model parameters

		Sex				
		Units	female	male	Combined	Years
Growth model	L_{∞}	cm			34.5	cm
	K				0.34	
	t_0				-0.143	
	Data source	Demestre <i>et al.</i> , 1997 (adopted by SGMED-08-03)				
Length weight relationship	a				0.0056	
	b				3.2488	
	M (scalar)	0.42				
	sex ratio (% females/total)	0.69				

L/W relationship from DCF; M from PRODBIOM

3 Fisheries information

3.1 Description of the fleet

Both species of red mullet, *Mullus surmuletus* and *M. barbatus*, are exploited by trawl and artisanal fleets in GSA 06, although small gears (trammel nets and gillnets) account only for 5% of the total landings of these species (Demestre et al., 1997). Trawl fisheries developed along the continental shelf and upper slope are multi-specific. Small vessels (12-16m length) operate mainly on the shallow shelf targeting on red mullets, octopus, cuttlefish and sea breams. Medium and large vessels usually operates on deep continental shelf and slope areas targeting on hake and decapod crustaceans, but some of these units can also operate on the shallow shelf depending on weather conditions or market prices. Red mullet is more intensively exploited from September to November coinciding with the recruitment period of this species (Martín et al., 1999). The total trawl fleet in the GSA 06 has declined from 810 boats in 1998 to 424 boats in 2016.

Table 3-1: Description of operational units exploiting the stock


	Country	GSA	Fleet Segment	Fishing Gear Class	Group of Target Species	Species
Operational Unit 1*	ESP	06	E – Trawl (12-24 m)	03 - Trawl	33-Demersal shelf species	Pagellus acarne Pagellus erythrinus Merluccius Octopus vulgaris Sepia officinalis Eledone cirrhosa

Table 3.1-2: Catch, bycatch, discards and effort by operational unit in the reference year

Operational Units*	Fleet (n° of boats)*	Catch (T or kg of the species assessed)	Other species caught (names and weight)	Discards (species assessed)	Discards (other species caught)	Effort (units)
[Operational Unit1]	424	1348 tons				48.6 Fishing days *1000
Total	424	1348 tons				48.6

3.2 Historical trends

The fishery developed in the early seventies and landings increased quickly. Since then landings widely fluctuates but a general decreasing trend is observed. In the period assessed landings fluctuate but without any clear trend. Fleet CPUE in the studied period fluctuates but no trend is observed.



3.3 Management regulations

- Engine power limited to 316 KW or 500 CV.
- Mesh size in the cod-end (50 mm diamond or 40 mm square).
- Fishing ban of trawl fishing in areas less than 50 m depth.
- Time at sea (12 hours per day and 5 days per week).
- Spatial and temporal closures of trawl fishing.
- Minimum legal size: 11 cm TL.

3.4 Reference points

Table 3.3-1: List of reference points and empirical reference values previously agreed (if any)

Indicator	Limit Reference point/emp irical reference value	Value	Target Reference point/empi rical reference value	Value	Comments
B		4210		3723	B mean as a referent point (B low = 2668)
SSB		1574		1287	SSB mean as a referent point (SSB low = 909)
F		0.74		0.26	F0.1 as a referent point
Y		1348		1115	Y mean as a referent point (Y low = 926)
CPUE		27.73		21.23	CPUE mean as a referent point (CPUE low = 16.78)

4 Fisheries independent information

4.1 {TYPE OF SURVEY}

Fill in one section for each of the direct methods used. The name of the section should be the name of the TYPE OF SURVEY.

4.1.1 Brief description of the direct method used

Description of the survey and method applied. One of several tables would have to be chosen: Egg Production Method, Acoustic survey, Trawl.

Direct methods: trawl based abundance indices

Table 4.1-1: Trawl survey basic information

Survey	MEDITS 2016	Trawler/RV	Miguel Oliver
Sampling season	SPRING		
Sampling design	random stratified with number of haul by stratum proportional to stratum surface		
Sampler (gear used)	GOC-73		
Cod –end mesh size as opening in mm	20		
Investigated depth range (m)	40-800		

Table 4.1-2: Trawl survey sampling area and number of hauls

Stratum	Total surface (km ²)	Trawlable surface (km ²)	Swept area (km ²)	Number of hauls
A (-50m)	3026			10
B (50-100m)	11314			36
C (100-200m)	6889			27
D (200-500 m)	6719			21
E (+500m)	4558			12
Total (km²)	32506			106

Map of hauls positions:




Table 4.1-3: Trawl survey abundance and biomass results

Depth Stratum	Years	kg per km ²	CV or other	N per km ²	CV or other
				
				
				
				
				
Total (... – ... m)				

*Comments

- Specify CV or other index of variability of mean
- Specify sampling design (for example random stratified with number of haul by stratum proportional to stratum surface; or systematic on transect;...)
- Specify if catchability coefficient is assumed =1 or other

Direct methods: trawl based length/age structure of population at sea

Slicing method

Report the maturity scale and age slicing method used

Table 4.1-4: Trawl survey results by length or age class

N (Total or sex combined) by Length or Age class	Year		

Total			

Sex ratio by Length or Age class	Year		

Total			

Comments

- ☐ Specify if numbers are per km² or raised to the area, assuming the same catchability .
- ☐ In case maturity ogive has not been estimated by year, report information for groups of years.
- ☐ Possibility to insert graphs and trends

Direct methods: trawl based Recruitment analysis

Table 4.1-5: Trawl surveys; recruitment analysis summary

Survey		Trawler/RV	
Survey season			
Cod –end mesh size as opening in mm			
Investigated depth range (m)			
Recruitment season and peak (months)			
Age at fishing-grounds recruitment			
Length at fishing-grounds recruitment			

Table 4.1-6: Trawl surveys; recruitment analysis results

Years	Area in km ²	N of recruit per km ²	CV or other

Comments

- Specify type of recruitment:
 - continuous and diffuse
 - discrete and diffuse □ discrete and localised
 - continuous and localised.
- Specify the method used to estimate recruit indices
- Specify if the area is the total or the swept one
- Possibility to insert graphs and trends

Direct methods: trawl based Spawner analysis

Table 4.1-7: Trawl surveys; spawners analysis summary

Survey		Trawler/RV	
Survey season			
Investigated depth range (m)			
Spawning season and peak (months)			


Table 4.1-8: Trawl surveys; spawners analysis results

Surveys	Area in km ²	N (N of individuals) of spawners per km ²	CV or other	SSB per km ²	CV or other

Comments


- Specify type of spawner:
 - total spawner
 - sequential spawner
 - presence of spawner aggregations
- Specify if the area is the total or the swept one
- Possibility to insert graphs e trends

4.1.2 Spatial distribution of the resources



4.1.3 Historical trends

MEDITS surveys data show a slight increasing trend in abundance along the period.



5 Ecological information

5.1 Protected species potentially affected by the fisheries

A list of protected species that can be potentially affected by the fishery should be incorporated here. This should also be completed with the potential effect and if available an associated value (e.g. bycatch of these species in T)

5.2 Environmental indexes

If any environmental index is used as i) a proxy for recruitment strength, ii) a proxy for carrying capacity, or any other index that is incorporated in the assessment, then it should be included here.

Other environmental indexes that are considered important for the fishery (e.g. Chl a or other that may affect catchability, etc.) can be reported here.

6 Stock Assessment

In this section there will be one subsection for each different model used, and also different model assumptions runs should be documented when all are presented as alternative assessment options.

6.1 Tunned VPA (XSA) (Darby & Flatman, 1994) and FLR libraries.

6.1.1 Model assumptions

6.1.2 Scripts

If a script is available which incorporates the stock assessment run (e.g. if using FLR in R) it should be provided here in order to create a library of scripts.

```
FLXSA.control.aa5 <- FLXSA.control(x=NULL, tol=1e-09, maxit=30, min.nse=0.3, fse=2.5,
rage=0, qage=2, shk.n=TRUE, shk.f=TRUE, shk.yrs=2, shk.ages=2,
window=100, tsrange=20, tspower=3, vpa=FALSE)
```

6.1.3 Input data and Parameters

Input parameters and model settings

Age group	M (Prodbiom)	Maturity (DCF)
0	1.23	0.127
1	0.41	0.929
2	0.28	0.999
3+	0.22	1.000

Growth parameters (Demestre et al, 1997)

$L_{inf} = 34.5$; $K = 0.34$; $T_0 = -0.143$

L/W relationship (DCF) a

$a = 0.0056$; $b = 3.2488$

- Proportion Fishing Mortality Prior to Spawning: 0.5
- Proportion Natural Mortality Prior to Spawning: 0.5

MODEL SETTINGS

XSA analysis. Catchability dependent on stock size for ages < 1

Catchability independent of age for ages ≥ 1

S. E. of the mean to which the estimates are shrunk = 2.5

Minimum standard error for population estimates = 0.3

6.1.4 Results


Fishing mortality at age:

	2004	2005	2006	2007	2008	2009	2010	2011
0	0.142	0.090	0.106	0.053	0.162	0.034	0.019	0.019
1	1.286	1.156	1.338	1.458	1.052	0.928	0.915	0.740
2	0.666	0.640	0.568	0.526	0.596	0.817	0.698	0.603
+gp	0.666	0.640	0.568	0.526	0.596	0.817	0.698	0.603

	2012	2013	2014	2015	2016
0	0.008	0.004	0.015	0.016	0.023
1	0.749	0.730	0.721	0.736	0.900
2	0.763	0.757	0.704	0.696	0.686
+gp	0.763	0.757	0.704	0.696	0.686

Other results:


Year	Recruits	SSB	F bar 1-2
2004	165995	909	0.976
2005	226478	1120	0.898
2006	181048	1391	0.953
2007	132283	1257	0.992
2008	124948	1031	0.824
2009	93792	1052	0.873
2010	113818	1102	0.806
2011	112569	1358	0.671
2012	114054	1391	0.756
2013	130860	1400	0.744
2014	128134	1498	0.713
2015	173039	1591	0.716
2016	125698	1633	0.793




Tables and graphs of Recruitment, SSB, Catch and F of the stock assessment model.

6.1.5 Robustness analysis


FLEET log catchability residuals



MEDITS log catchability residuals



6.1.6 Retrospective analysis, comparison between model runs, sensitivity analysis, etc.



Retrospective analysis performed with XSA of Recruitment, SSB and Harvest.

6.1.7 Assessment quality

Stability of the assessment, evaluation of quality of the data and reliability of model assumptions.

6.2 Y/R (version 3.1.1; NOAA Fisheries Tools)

Model parameters

Age group	Stock weight	Catch weight	SSB weight	Maturity	M	Selectivity
0	0.011	0.011	0.011	0.127	1.23	0.02
1	0.035	0.035	0.035	0.929	0.41	0.54
2	0.087	0.087	0.087	0.999	0.28	1.00
3+	0.161	0.161	0.161	1.000	0.22	1.00

First Age in Data: 0

Last Age in Data: 5

Age in Plus Group: 3


Y/R results

F 0.1 0.26

F max 0.47

F current (*) 0.74

(*) From XSA. Mean \bar{F} 1-2 in 2014-2016.



6.3 STOCK / RECRUITMENT RELATIONSHIP

7 Stock predictions

When an analytical assessment exists, predictions should be attempted. All scenarios tested (recruitment and/or fishing mortality) should be reported. The source of information/model used to predict recruitment should be documented.

7.1 Short term predictions

7.2 Medium term predictions

7.3 Long term predictions

8 Draft scientific advice

Based on	Indicator	Analytic al reference point (name and value)	Current value from the analysis (name and value)	Empirical reference value (name and value)	Trend (time period)	Stock Status
Fishing mortality	Fishing mortality	$F_{0.1}$	0.26	$F_{current}$ (ages 1-2) = 0.74	D	O_H
	Catch		1348 (2016)	Mean catch (2004-2016) = 1115 tons	N	
Stock abundance	Total Biomass		4210 (2014- 2016)		N	O_H
	SSB		1574 (2014- 2016)	33 th percentile = 1120 66 th percentile =1391	I	
Recruitment		126×10^6 N	(in 2016)			
Final Diagnosis		In High overfishing ($F_{current} > F_{0.1}$). Relative high total SSB				

State the rationale behind that diagnoses, explaining if it is based on analytical or on empirical references

8.1 Explanation of codes

Trend categories

- 1) N - No trend
- 2) I - Increasing
- 3) D – Decreasing
- 4) C - Cyclic

Stock Status

Based on Fishing mortality related indicators

- 1) **N - Not known or uncertain** – Not much information is available to make a judgment;
- 2) **U - undeveloped or new fishery** - Believed to have a significant potential for expansion in total production;
- 3) **S - Sustainable exploitation**- fishing mortality or effort below an agreed fishing mortality or effort based Reference Point;
- 4) **IO –In Overfishing status**– fishing mortality or effort above the value of the agreed fishing mortality or effort based Reference Point. An agreed range of overfishing levels is provided;

Range of Overfishing levels based on fishery reference points

In order to assess the level of overfishing status when $F_{0.1}$ from a Y/R model is used as LRP, the following operational approach is proposed:

- If $F_c/F_{0.1}$ is below or equal to 1.33 the stock is in (**O_L**): **Low overfishing**
 - If the $F_c/F_{0.1}$ is between 1.33 and 1.66 the stock is in (**O_I**): **Intermediate overfishing**
 - If the $F_c/F_{0.1}$ is equal or above to 1.66 the stock is in (**O_H**): **High overfishing**
- * F_c is current level of F

- 5) **C- Collapsed**- no or very few catches;

Based on Stock related indicators

- 1) **N - Not known or uncertain**: Not much information is available to make a judgment
- 2) **S - Sustainably exploited**: Standing stock above an agreed biomass based Reference Point;
- 3) **O - Overexploited**: Standing stock below the value of the agreed biomass based Reference Point. An agreed range of overexploited status is provided;

Empirical Reference framework for the relative level of stock biomass index

- **Relative low biomass**: Values lower than or equal to 33rd percentile of biomass index in the time series (**O_L**)
- **Relative intermediate biomass**: Values falling within this limit and 66th percentile (**O_I**)

- **Relative high biomass:** Values higher than the 66th percentile (**O_H**)

- 4) **D – Depleted:** Standing stock is at lowest historical levels, irrespective of the amount of fishing effort exerted;
- 5) **R –Recovering:** Biomass are increasing after having been depleted from a previous period;

Agreed definitions as per SAC Glossary

Overfished (or overexploited) - A stock is considered to be overfished when its abundance is below an agreed biomass based reference target point, like *B_{0.1}* or *B_{MSY}*. To apply this denomination, it should be assumed that the current state of the stock (in biomass) arises from the application of excessive fishing pressure in previous years. This classification is independent of the current level of fishing mortality.

Stock subjected to overfishing (or overexploitation) - A stock is subjected to overfishing if the fishing mortality applied to it exceeds the one it can sustainably stand, for a longer period. In other words, the current fishing mortality exceeds the fishing mortality that, if applied during a long period, under stable conditions, would lead the stock abundance to the reference point of the target abundance (either in terms of biomass or numbers)